

# SULPHASOD

## Injection

For Veterinary Use Only

### COMPOSITION:

Each ml contains  
Sulphadimidine Sodium.....333mg

### SPECTRUM OF ACTIVITY:

Sulfdimidine Sodium is a sulphonamide with antimicrobial activity against Gram-positive cocci (hemolytic streptococci, pneumococci and some staphylococci), against some Gram-negative bacteria (like Pasteurella spp., E. coli, Salmonella spp.) and against some Rickettsiae and protozoa. Sulfdimidine is rapidly absorbed from the site of application and slowly excreted by the organism, so in most animals a single daily dose is sufficient. Bacterial resistance against sulphonamides mostly is permanent and cross-resistance between sulphonamides is common.

### INDICATIONS:

**Sulphasod** Injection is indicated for pneumonia, foot rot, diphtheria in calves, Actinobacillosis, Colibacillosis, Salmonellosis, Pasteurellosis (Haemorrhagic septicemia), enteritis, poly arthritis, Streptococcal mastitis, dysentery, lymphangitis and Coccidiosis in sheep, goats, cattle and calves. For treatment of respiratory, gastro-intestinal infections, cholera and coccidiosis in poultry.

### Side-effects:

Sometimes hemorrhagic enteritis (in cattle), kidney damage, hypersensitivity reactions or crystalluria may occur.

### DOSAGE AND ADMINISTRATION:

For intravenous or deep intramuscular injection.

**Cattle and Horses:** 30ml per 50 kg bodyweight followed by 15ml per 50 kg bodyweight for 2-5 days.

**Sheep and goat:** 15ml per 25kg bodyweight for 2-5 days.

**Dogs/Cats and other Small Animals:**

3.0 ml per 5 kg bodyweight for 2-5 days.

**Poultry:** 2.5ml per litre of drinking water for 3 days...

### CONTRAINDICATIONS:

Hypersensitivity to sulfonamides.

Administration to animals with a seriously impaired renal and/or hepatic function or with blood dyscrasias.

### ADVERSE REACTION:

Sulfonamide or its metabolites can precipitate in urine, and it is easier to produce crystallization at high doses and long-term administration, causing crystalluria, hematuria, leucopenia or blockage of renal tubules.

### WITHDRAWAL TIME:

**Meat:** 14 days.


**Milk:** 6 days.

### PRECAUTIONS:

Store between 15-25° C in a cool and dry place. Gently warm the bottle in water bath if crystals appear in the solution.

Keep out of the reach of children. Consult the veterinarian before use.

### BP Vet. Specs.



 **Selmore**  
Pharmaceuticals (Pvt) Ltd.

Manufactured by:  
Selmore Pharmaceuticals (Pvt) Ltd.  
36 Km Multan Road, Lahore, Pakistan.



ISO CERTIFIED COMPANY