CALPHO-M Injection

For Veterinary Use Only

COMPOSITON:

Each ml contains	
Calcium Gluconate	208.30mg
Magnesium Hypophosphite	53.30mg
Magnesium Chloride	
Calcium D Saccharate	
Boric Acid	43.30mg
Dextrose	200mg

INDICATIONS:

Calpho-M injection is indicated for the treatment of hypocalcemia, complicated by magnesium deficiency and hypoglycemia, pregnancy toxemia and milk fever in cattle and dairy animals

PROPERTIES:

Calpho-M injection contains active substances that can quickly restore calcium, magnesium and energy balance. Calcium is essential for bone-formation, normal functioning of the nervous and muscular systems and regulates acid-alkaline balance. Calcium gluconate is a highly tolerated calcium source able to rapidly replenish the levels of this mineral in the body. It reduces blood vessel permeability and fragility, moderates muscular and neural excitability and has tonic action on the heart.

Magnesium hypophosphite and magnesium chloride provides highly bioavailable magnesium, which exerts important biological functions, in proper ratio with calcium for maintenance of normal neuromuscular excitability.

Boric acid has a stabilizing action; it eliminates side effects and allows, besides the intravenous. also subcutaneous, intramuscular and intraperitoneal administration. While Dextrose is a rich source of energy.

DOSAGE AND ADMINISTRATION:

Administer the following dose by subcutaneous or slow intravenous injection only. Cattle/Horse: 200-300 ml Sheep/Goat: 30-60 ml Additional repeat treatments can be given at 24hour interval

PRECAUTIONS:

- The solution should be warmed to body temperature before administration.
- Intravenous administration must be carried out very slowly over a period of 30 minutes.
- Heart rate and respiration must be continuously monitored. In case of cardiac arrhythmia, fall in blood pressure and agitation, infusion must be stopped immediately.
- Do not injection Calpho-M along with other medicinal products like penicillin, glucocorticoids, steroids, menbutone, sodium phosphate, promethazine HCI, sodium bicarbonate, streptomycin, tetracycline and sulpha salts is strongly prohibited.
- Store between 15-25°C in a cool and dry place.
- Consult the veterinarian before use.
- · Keep out of the reach of children.

Selmore's Specs.





Selmore Pharmaceuticals (Pvt) Ltd.

