

CAMILOX

INJECTION

For Veterinary Use Only

COMPOSITION:

Each ml contains:

Meloxicam7.5mg

INDICATIONS:

CamiloX Injection is indicated for the treatment of acute or chronic pain, fever inflammation, musculoskeletal disorders, arthritis, tenosynovitis, bursitis and postoperative pain and inflammation.

Cattle/sheep/goat: For use in acute respiratory infection with appropriate antibiotic therapy to reduce clinical signs. For use in diarrhea in combination with oral rehydration therapy to reduce clinical signs in calves of over one week of age and young, non-lactating cattle.

Dogs: Alleviation of inflammation and pain in both acute and chronic musculo-skeletal disorders. Reduction of post-operative pain and inflammation following orthopedic and soft tissue surgery.

Cats: Reduction of post-operative pain after ovariohysterectomy and minor soft tissue surgery.

CONTRAINDICATIONS:

Cattle

- Do not use in animals suffering from impaired hepatic, cardiac or renal function and haemorrhagic disorders, or where there is evidence of ulcerogenic gastrointestinal lesions.
- Do not use in case of hypersensitivity to the active substance or to any of the excipients.
- For the treatment of diarrhoea in cattle, do not use in animals of less than one week of age.

Dogs and cats

- Do not use in pregnant or lactating animals.
- Do not use in animals suffering from gastrointestinal disorders such as irritation and haemorrhage, impaired hepatic, cardiac or renal function and haemorrhagic disorders.
- Do not use in case of hypersensitivity to the active substance or to any of the excipients.
- Do not use in animals less than 6 weeks of age nor in cats of less than 2 kg.

ADVERSE REACTIONS:

Cattle

Subcutaneous, intramuscular as well as intravenous administration is well tolerated; only a slight transient swelling at the injection site following subcutaneous administration was observed in less than 10% of the cattle treated in clinical studies. In very rare cases anaphylactoid reactions which may be serious (including fatal) may occur and should be treated symptomatically.

Dogs and cats

Typical adverse reactions of NSAIDs such as loss of appetite, vomiting, diarrhoea, faecal occult blood, lethargy and renal failure have occasionally been reported. In very rare cases elevated liver

enzymes have been reported.

In dogs, in very rare cases, haemorrhagic diarrhoea, haematemesis, and gastrointestinal ulceration have been reported.

In very rare cases anaphylactoid reactions may occur and should be treated symptomatically.

If adverse reactions occur, treatment should be discontinued and the advice of a veterinarian should be sought.

TARGET SPECIES:

Cattle, horse, sheep, goat, dog and cats.

DOSAGE & ADMINISTRATION:

Administer the following dose intravenously (IV), subcutaneously (SC) or intramuscularly (IM) at 0.2mg/kg body weight.

Cattle/Horses: 6-8ml per 200kg body weight.

Sheep/Goat: 0.5-1ml per 20-35kg body weight.

Dog/Cat: 0.2-0.5ml for 5-20kg body weight.

PRECAUTION

Do not use in cattle producing milk for human consumption, including non-lactating dairy cows during the dry period.

Keep out of the reach of Children.

Consult the veterinarian before use.

STORAGE CONDITIONS AND SHELF LIFE


Store between 15-25°C in a cool and dry place.

Shelf Life: 2 years.

PACKING

50ml

BP Vet Specs



 **Selmore**
Pharmaceuticals (Pvt) Ltd.

Manufactured by:
Selmore Pharmaceuticals (Pvt) Ltd.
36 Km Multan Road, Lahore, Pakistan.

