

1g CEFTISEL

DRY POWDER INJECTION

For Veterinary Use Only

COMPOSITION:

Each vial contains:

Ceftiofur Sodium equivalent to Ceftiofur.....1g

PHARMACEUTICAL FORM:

Powder and solvent for solution for injection

PHARMACOLOGICAL PROPERTIES:

Ceftisel Dry Powder Injection contains the sodium salt of Ceftiofur which is a third-generation, broad spectrum cephalosporin antibiotic active against gram-positive and gram-negative bacteria including β -lactamase-producing strains. Like other cephalosporin, Ceftiofur is bactericidal in vitro, resulting from inhibition of cell wall synthesis. It is rapidly metabolized to desfuroyl Ceftiofur, which accounts for most of its antimicrobial activity. This product may be used in lactating dairy cattle, sheep, and goats.

INDICATIONS:

Cattle/ Buffalo: CEFTISEL injection is indicated for the treatment of cattle with acute bacterial respiratory disease in which Mannheimia haemolytica, Pasteurella multocida or Histophilus somni are involved.

Sheep/Goats: CEFTISEL injection is indicated for the treatment of respiratory disease (pneumonia) associated with Mannheimia haemolytica and Pasteurella multocida.

Horses: CEFTISEL injection is indicated for the treatment of bacterial respiratory disease in which Streptococcus spp including Streptococcus zooepidemicus and Streptococcus equi, Staphylococcus spp and/or Pasteurella spp are involved.

DOSAGE & ADMINISTRATION:

Cattle: (use intramuscularly) 1ml per 50 kg bodyweight.

Sheep/Goat: (use intramuscularly) 2ml per 50kg bodyweight.

Horses: (use intramuscularly) 2ml per 50kg bodyweight.

Dog: (use subcutaneous route) 0.2ml per 5kg body weight.

Day old chicks: (use sub cutaneous route) 1ml per 250 to 625 day old chicks

CONTRAINDICATIONS:

- Do not use in animals previously found to be hypersensitive to Ceftiofur and other lactam antibiotics?
- Do not use in case of known resistance to the active substance.
- Do not use in cases where resistance to other cephalosporin or beta-lactam antibiotics has occurred.
- Do not use in poultry (including eggs) due to risk of spread of antimicrobial resistance to humans.

Adverse reactions (frequency and seriousness)

General symptoms are not detected. The use of ceftiofur sodium may result in some signs of immediate and a short lasting pain at the site of injection.

PRECAUTIONS:

Consult the veterinarian before use.


Keep out of the reach of children.

Store between 15-25 °C in a cool and dry place.

WITHDRAWAL TIME:

Meat: 1 days.

BP SpecsOSITION:



Manufactured by:
Selmores Pharmaceuticals (Pvt) Ltd.
36 Km Multan Road, Lahore, Pakistan.

