

COLIMOXIN

Injection

For Veterinary Use Only

STATEMENT OF ACTIVE SUBSTANCE (S) AND OTHER

INGREDIENTS:

Each ml contains:

Amoxicillin as Trihydrate.....100mg
Colistin Sulphate.....250,000IU

INDICATIONS:

Colimoxin Injection is indicated for the treatment of infections caused by gram-positive and gram-negative microorganisms of respiratory tract (bronchial and lungs infections), gastrointestinal tract, urogenital tract and cutaneous systems.

CONTRAINDICATIONS:

Not for intravenous administration. Do not use in animals with known hypersensitivity to the active ingredients.

ADVERSE REACTIONS:

Occasional local reaction may occur. Occasional allergies to the penicillin have been observed but these are rare.

TARGET SPECIES:

Cattle, sheep

DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION:

Administer the following dose once or twice daily by deep intramuscular injection. 1ml per 10kg body weight per day.

Advice on correct administration:

To ensure a correct dosage body weight should be determined as accurately as possible.

WITHDRAWAL PERIOD (S):

Meat: 21 days

Milk: 3 days

PRECAUTIONS:

Store between 15-25°C in a cool and dry place.

Shake well before use.

Keep out of the reach of children.

Consult the veterinarian before use only

Protect from light. Do not use this veterinary medicinal product after the expiry date which is stated on the carton.

SPECIAL WARNINGS FOR EACH TARGET SPECIES:

Massage the injection site after administration. In adult cattle, the volume should be divided between two injection sites.

SPECIAL PRECAUTIONS FOR USE IN ANIMALS:

Use of the product should be based on susceptibility testing of the bacteria isolated from the animal. If this is not possible, therapy should be based on local (regional, farm level) epidemiological information about susceptibility of the target bacteria.

Special precautions to be taken by the person administering the veterinary medicinal product to animals: Penicillin and cephalosporins may cause hypersensitivity (allergy) following

injection, inhalation, ingestion or skin contact. Hypersensitivity to penicillin may lead to cross sensitivity to cephalosporins and vice versa. Allergic reaction to these substances can occasionally be serious.

1. Do not handle this product if you know you are sensitised or if you have been advised not to work with such preparations.
2. Handle this product with great care to avoid exposure, taking all recommended precautions.
3. If you develop symptoms following exposure, such as a skin rash, you should seek medical advice and show the doctor this warning. Swelling of the face, lips and eyes or difficulty with breathing are more serious symptoms and require urgent medical attention.

PREGNANCY AND LACTATION:

Amoxicillin is safe for use in pregnant animals.

INTERACTION WITH OTHER MEDICINAL PRODUCTS AND OTHER FORMS OF INTERACTION:

Tetracyclines are bacteriostatic antibiotics that presumably may interfere with a bactericidal agent such as amoxicillin. Since Amoxicillin acts by inhibiting cell wall synthesis, agents such as tetracyclines, which inhibit protein synthesis, could mask the bactericidal effect of amoxicillin.

OVERDOSE (SYMPTOMS, EMERGENCY PROCEDURES, ANTIDOTES):

Do not exceed the stated dose. Hypersensitivity (allergic) reactions to penicillins can vary from localised swelling to anaphylaxis and death. Therapy involves hot- or cold-water soaks and/or corticosteroids.


INCOMPATIBILITIES:

None known.

SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS IF ANY:

Any unused product or waste material should be disposed of in accordance with national requirements.

Innovator's Specs.



Selmores
Pharmaceuticals (Pvt) Ltd.

Manufactured by:
Selmores Pharmaceuticals (Pvt) Ltd.
36 Km Multan Road, Lahore, Pakistan.



ISO CERTIFIED COMPANY