

# 120x189mm

## Quina-CS

2.5gm

INJECTION



**For Veterinary Use Only**

### COMPOSITION:

Each 2.5gm vial contains

Quinapyramine Sulphate (B.Vet. C) ..... 1.5gm

Quinapyramine Chloride (B.Vet. C) ..... 1gm

### Pharmaceutical Form:

Powder for Injection

### INDICATIONS:

QUINA-CS Injection is Quinapyramine Sulphate mixed with Quinapyramine Chloride active against Trypanosomabrucei, T. Congolenses, T. equinum, T. equiperdum, T. evansi, T. simiae and T. vivax in various species of animal viz. cattle, buffalo, horses, camels, donkey and dogs.

### CONTRAINDICATIONS:

QUINA-CS is contraindicated in animals previously found to be hypersensitive to the drug. The drug should not be used in young stock because it cause sweating, salivation, polypnoea, tachycardia and death might occur.

### ADVERSE REACTIONS:

Transient salivation, restlessness and muscular tremors may be observed in some weak animals.

### TARGET SPECIES:

Cattle, buffalo, horses, camels, donkey and dogs.

### DOSAGE & ADMINISTRATION:

Add 15 ml distilled water into the vial; shake well to make suspension for immediate use. Administer by subcutaneous or intramuscular injection. Massage the injection site to disperse the drug and minimize the risk of local reaction. In endemic areas of Surra and Nagana the animals should be injected at intervals of 2 - 3 months as a prophylactic measure again.

Recommended dose is 2ml per 45kg body weight by SC or IM injection. Divide more than 15ml dose in 2-3 different injection sites.

### WITHDRAWAL PERIOD:

Meat: 21 days

Milk: 4 days

### STORAGE CONDITIONS AND SHELF LIFE:

Store below 25°C without freezing and refrigerating.

Keep away from sunlight.

Do not store the reconstituted suspension.

### SPECIAL WARNINGS FOR EACH TARGET SPECIES:

In horses local reactions to S.C.I. may occur at the sites of injection so the dose must be divided into 2 or 3 injections at different sites or inject I.M.

The prepared suspension for immediate use, and shake well before use.

Estimate live weight carefully and allow animals' access to shade and water after dosing.

Massage the injection site well.

The dose may be divided into 2 doses with 12 hours intervals.

Estimate live weight carefully and allow animals access to shade and water after dosing.

Do not store the reconstituted suspension.

### SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED VETERINARY MEDICINAL PRODUCT OR WASTE MATERIALS DERIVED FROM THE USE OF SUCH PRODUCTS:

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements..

### PACKING:

2.5gm/vial



Manufactured by:  
Selmore Pharmaceuticals (Pvt) Ltd.  
36-Km Multan Road, Lahore Pakistan.

