

DYRO-X

Powder

For Veterinary Use Only

Composition:

Each 12gm contains:	
Neomycin Sulphate.....	400mg
Streptomycin Sulphate.....	400mg
Sulfaguanidine.....	4gm
Kaolin.....	4gm
Pectin.....	400mg
Bismuth Subnitrate.....	2gm
Vitamin A Acetate.....	800,000IU

Neomycin Sulfate:

Neomycin is an antibiotic that fights bacteria in the animal's body. Suppression of Intestinal Bacteria: Neomycin sulfate are indicated as adjunctive therapy as part of a regimen for the suppression of the normal bacterial flora of the bowel, eg, preoperative preparation of the bowel.

Streptomycin Sulphate:

Streptomycin Sulfate is the sulfate salt form of streptomycin, an aminoglycoside antibiotic derived from *Streptomyces griseus* with antibacterial property. Streptomycin is indicated for the treatment of individuals with moderate to severe infections caused by susceptible strains of microorganisms.

Sulfaguanidine:

Sulfaguanidine is a guanidine derivative of sulfanilamide used in veterinary medicine. Sulfaguanidine is poorly absorbed from the gut but is well suited for the treatment of bacillary dysentery and other enteric infections.

Kaolin is a clay mineral, part of the group of industrial minerals. It is a layered silicate mineral. Kaolin is used for mild-to-moderate diarrhea, severe diarrhea (dysentery), and cholera.

Pectin: Pectin is a naturally occurring substance (a polysaccharide) found in berries, apples and other fruit...In the pharmaceutical industry, it is used to reduce blood cholesterol levels and gastrointestinal disorders

Bismuthsubnitrate: also referred to as bismuth oxynitrate or bismuthyl nitrate, is a highly water-soluble crystalline compound that has been used as a treatment for duodenal ulcers and anti-diarrheic agent.

Vitamin A: is present in foods such as retinol and

retinyl esters (usually retinyl acetate) aids in digestion process along with other uses.

Indications:

Dyro-X Powder is indicated for the treatment of diarrhea, indigestion, impaction, bacterial enteritis and other digestive disorders due to susceptible microorganism in domestic animals.

Dosage and Administration:

Administer 12g in drinking water or feeding per 40kg weight of domestic animals. Repeat the dose after 12 hours if required.

Precautions:

Consult the veterinarian before use. Protect from light & moisture.

Keep out of the reach of children. Store between 15-25°C in a cool and dry place.

Withdrawal Time:

Animals must not be slaughtered for human consumption before 7 days from the last treatment.

Innovator's Specs.



Manufactured by:
Selmores Pharmaceuticals (Pvt) Ltd.
36 Km Multan Road, Lahore, Pakistan.



ISO CERTIFIED COMPANY