FON-BIO

ORAL POWDER

For Veterinary Use Only

COMPOSITION:

Each Gram contains:

Neomycin Sulphate	.150mg
Florfenicol	100mg
Oxytetracycline HCI	. 300mg

INDICATIONS:

FON-Bio oral Powder is indicated in Gastrointestinal, upper and lower respiratory tract infections. E-coli and Salmonella.

Fon-Bio Powder is indicated for the prevention and treatment of gastrointestinal, digestive and urogenital infections caused by these antibiotic sensitive micro-organism like salmonella, Pasteurella, Haemophilus, E.coli, mycoplasma spp. in poultry, calves, sheep & goats.

DESCRIPTION:

Fon-Bio Powder is a combination of 3 broad spectrum antibiotics. Neomycin belongs to a class of drugs known as amino glycosides antibiotic. Florfenicol is readily absorbed from GIT and actively penetrates into the majority of organs and tissues of organism. Oxytetracycline works by interfering with the ability of bacteria to produce essential proteins without those proteins, the bacteria cannot grow multiply and increase in numbers. Show excellent results against typhoid, mycoplasma, E.coli, Pasteurella, Haemophilus, streptococcus & staphylococcus spp.

MECHANISM OF ACTION:

Neomycin is an aminoglycoside broad-spectrum bactericidal antibiotic with activity against certain members of the Enterobacteriaceae. Its mode of action is at the ribosomal level. Neomycin sulphate is recommended for the treatment of infections of the alimentary tract, caused by Gram-positive and Gram-negative bacteria that are sensitive to neomycin. Oxytetracycline HCl is a broad-spe ctrum antibiotic effective against Actinomycosis, Rickettsia, Mycoplasma and Chlamydia spp. and is therefore recommended in the treatment of Pasteurella multocida and Mycoplasma infections in poultry. Florfenicol is a broad-spectrum bacteriostatic antibiotic effective against Staphylococcus, Streptococcus, Pneumococcus, Corynebacterium spp., Coli bacteria. Klebsiella spp. Typhus and Para typhus bacteria, Shigella spp., Brucella spp., Haemophilus spp., Pasteurella spp., Vibrio spp. and Bacillus anthraces.

DOSAGE & ADMINISTRATION:

Administer 1 gram in 4 Liters of drinking water for 3-5 days.

PRECUATIONS:

Keep the Jar / Sachet tightly closed after initial use. Consult the veterinarian before use.

Protect from light & moisture.

Keep out of the reach of children.

Store between 15-25 °C in a cool and dry place.

WITHDRAWAI TIME.

Meat and Eggs: 07 Days

Innovator's Specs.





