Helmex

For Veterinary Use Only

COMPOSITION

Each ml contains Levamisole HCI15mg Oxyclozanide30mg Cobalt Sulphate1.67mg Sodium Selenite0.50mg

INDICATIONS

Helmex plus Drench is effective in cattle, sheep. goat for all types of gastrointestinal worms including Ostertagia and Nematodirus, Lung worms, tape worms segments and liver flukes. It covers the deficiency of cobalt sulphate and sodium selenite.

Helmex plus Drench has a high safety margin due to which it equally effective at all ages including young, pregnant and lactating animals with no diet restriction.

PHARMACODYNAMICS

Levamisole acts on the acetylcholine receptors of worms leading to paralysis and death of parasite. Levamisole also restores the body immune system by chemo taxis of macrophages and T-lymphocyte. Oxyclozanide belongs to Salicylanilide group of anthelmintic which is effective against Fasciola species, acting as an uncouple of oxidative phosphorylation of parasites. It may act at more than one site to decrease the levels of ATP leading to metabolic malfunction and death of parasite.

Cobalt helps in the synthesis of vitamin B-12 in ruminants, while selenium acts as an antioxidant and works against the oxidizing effects of parasitic toxins results relief from parasitic stress.

PHARMACOKINETICS

Levamisole is quickly absorbed into blood stream after oral administration. It is quickly distributed into throughout the whole body, including body fluids like bronchial mucus and tears. Levamisole is metabolized in liver and after 24 hours of administration 90% dose is excreted through kidnevs.

Oxyclozanide is slowly absorbed after oral administration with peak plasma levels 24 hours after dosing. Excretion is predominantly fecal, biliary excretion being the most important route of elimination (cattle studies only).

DOSAGE AND ADMINISTRATION

The normal dosage is 5ml per 10Kg body weight.

| Cattle | | Sheep / Goat | |
|----------------------|---------|-----------------------|--------|
| 1ml / 2kg bodyweight | | 1ml / 2 kg bodyweight | |
| Bodyweight | Dose | Bodyweight | Dose |
| 75kg | 37.5ml | 15kg | 7.5ml |
| 76-125kg | 62.5ml | 16-25kg | 12.5ml |
| 126-175kg | 87.5ml | 26-35kg | 17.5ml |
| 176-225kg | 112.5ml | 36-45kg | 22.5ml |
| 225-300kg | 150ml | 66-85kg | 42.5ml |

Withdrawal Period Meat: 14 days, Milk: 24 hours

PRECAUTIONS

- Shake well before use.
- Store between 15-25°C in a cool and dry place.
- Consult the veterinarian before use
- Keep out of the reach of Children.

Innovator's Specs





