

# NOVAMOX-LA

## INJECTION

For Veterinary Use Only

### COMPOSITION:

Each ml contains  
Amoxicillin as Trihydrate.....150mg.

### INTRODUCTION:

Novamox-LA Injection is a broad spectrum semi-synthetic penicillin, bactericidal in action. In vitro it is effective against a wide range of gram positive and gram negative pathogenic bacteria which include, Actinobacillus equi, Actinobacillus lingers, Actinomyces bovis, Bacillus anthracis, Bordetella bronchiseptica, Clostridium species, Corynebacterium species, Erysipelothrix rhusiopathiae, Escherichia coli, Fusiformis species, Haemophilus species, Moraxella species, Pasteurella species, Proteus mirabilis, Salmonella species, Staphylococci and Streptococci in cattle, sheep, dogs and cats.

### INDICATIONS:

For the treatment of upper respiratory tract, gastrointestinal tract, urogenital tract infections, skin and soft tissues infection and in prevention of post-operative infections. It is a drug of choice for the treatment of pneumonia, ear, throat, enteritis, E.coli, wound infections, mastitis, metritis. In poultry for coryza, enteritis, streptococcus's and other respiratory diseases.

### DOSAGE & ADMINISTRATION:

**Cattles and Sheep:** by intramuscular injection only.

**Dogs and Cats:** by subcutaneous or intramuscular injection. Poultry: by subcutaneous or intramuscular injection.

**Cattle, Sheep, Dogs, Horses:**

1ml for 10kg body weight, repeat after 48 hours if required.

**Poultry:** 0.25ml per 1kg body weight.

Normal aseptic precautions should be observed, if dose volume exceeds 20 ml in cattle, It should be divided and injected into two sites.

### CONTRAINDICATIONS:

This product is not suitable for administration via intravenous or intrathecal routes.

Milk for human consumption must not be taken from a cow during treatment. Milk for human consumption may only be taken after 24 hours from the last treatment. Not for use in sheep producing milk for human consumption. Animals must not be slaughtered for human consumption during treatment. Cattle may be slaughtered for human consumption only after 14 days from the last treatment Sheep may be slaughtered for human consumption only after 10 days from the last treatment.

Occasional local tissue reaction may result from use of this product as with other penicillins, Amoxicillin should not be used in rabbits, hamsters.

### OPERATOR WARNINGS:

Penicillins and cephalosporin may cause hypersensitivity (allergy) following injection, Inhalation, Ingestion or skin contact. Hypersensitivity to penicillin may lead to cross reactions to cephalosporin and vice versa. Allergic reactions to these substances may occasionally be serious.

1. Do not handle this product if you know you are sensitized, or if you have been advised not to work with such preparations.

2. Handle this product with great care to avoid exposure taking all recommended precautions.

3. If you develop symptoms following exposure, such as a skin rash, you should seek medical advice and show the doctor this warning. Swelling of the face, lips or eyes or difficulty breathing are more serious symptoms and require urgent medical attention.

### PRECAUTIONS:

Shake well before use.

Store between 15-25°C in a cool and dry place.

Do not use in animals sensitive to penicillin


Keep out of the reach of children.

Consult the veterinarian before use

### WITHDRAWAL TIME:

Meat: 28 days.

**BP Vet. Specs.**



 **Selmores**  
Pharmaceuticals (Pvt) Ltd.

Manufactured by:  
Selmores Pharmaceuticals (Pvt) Ltd.  
36 Km Multan Road, Lahore, Pakistan.

