

ODEMTIC

50mg



For Veterinary Use Only

Injection

Statement of active substance (s) and other ingredients:

Each ml contains
Furosemide (as monoethanolamine)50mg

Indications:

Odemtic 50mg Injection treats congestive heart failure, pulmonary edema (fluid in the lungs), high blood pressure, high calcium and potassium levels, liver disease and some kidney diseases. It is a potent diuretic drug used in animals who suffer from fluid retention.

Contraindications:

Do not use in cases of liver coma, renal deficiency with anuria, electrolyte deficiency (hypokalemia, hyponatremia), hypotony and sulfonamide allergy.

Adverse reactions:

Odemtic 50mg Injection is a highly effective diuretic and, if given in excessive amounts, as with any diuretic, may lead to excessive diuresis which could result in electrolyte imbalance, dehydration and reduction of plasma volume, enhancing the risk of circulatory collapse, thrombosis and embolism.

The concurrent use of furosemide with some antibiotics may be inadvisable. There is evidence that the drug enhances the nephrotoxic potential of aminoglycosides, cephalosporins and polymyxins and increases the ototoxic effects of all aminoglycosides.

In cases of longer therapy, hypokalemia and hyponatremia may occur.

Too rapid injection in dogs may cause staggering and vomiting.

Use during pregnancy, lactation or lay:

Can be used during pregnancy and lactation.

Target species:

Cattle, buffalo, horses, camel, sheep, goat, dogs and cats.

Dosage for each species, route(s) and method of administration:

Administer the following dose by IM or IV injection once or twice daily until desired results are achieved.

Cattle/Buffalo: Inject intramuscularly or intravenously 10ml once daily or 5ml twice daily at 12 hours interval.

Sheep/Goat: Inject intramuscularly or intravenously 1ml per 50 kg body weight.

Horses/Camel: Inject 5 to 10ml intramuscularly or intravenously once or twice daily at 06-08 hours interval.

Dog/Cats: Administer intravenously or intramuscularly 0.25-0.5 ml per 5 kg body weight once or twice daily at 06-08 hours interval.

During prolonged treatment, it is necessary to monitor the serum electrolyte balance.

Overdose (symptoms, emergency procedures, antidotes), if necessary:

Doses higher than recommended may cause transitory deafness. Cardiovascular side effects may be observed in weak and old patients following overdosage.

Withdrawal period(s):

Meat: 02 days.

Milk: 02 days.

Storage precautions:

Odemtic 50mg Injection is a highly effective diuretic-saluretic which if given in excessive amounts may result in dehydration and electrolyte imbalance.

Concurrent use with aminoglycoside antibiotics may increase risks of nephrotoxicity and ototoxicity.

Odemtic 50mg Injection is contraindicated in animals with anuria or who are hypersensitive to the drug. Administration of NSAIDs with furosemide may diminish the effect.

Consult the veterinarian before use.

Store between 15-25°C in a cool and dry place.

Keep out of the reach of children.

Keep container in the outer carton in order to protect from light.

Special precautions for the disposal of unused product or waste materials if any:

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements.

BP Specs.



Manufactured by:
Selmore Pharmaceuticals (Pvt) Ltd.
36-Km Multan Road, Lahore Pakistan.
Head Office:
82, R-1, MA Johar Town, Postal Code 54100
Lahore, Pakistan.



ISO CERTIFIED COMPANY