

# 118x188mm

## SELMOX LA INJECTION

For Veterinary Use Only

### COMPOSITION:

Each ml contains  
Amoxicillin (as Trihydrate).....200mg

### PRODUCT USES:

Selmox LA Injection is a broad spectrum semi-synthetic penicillin, bactericidal in action. In vitro it is effective against a wide range of Gram-positive and Gram-negative pathogenic bacteria which include: Actinobacillus equuli, Actinobacillus lignieresii, Actinomyces bovis, Bacillus anthracis, Bordetella bronchiseptica, Clostridium species, Corynebacterium species, Erysipelothrix rhusiopathiae, Escherichia coli, Fusiformis species, Haemophilus species, Moraxella species, Pasteurella species, Proteus mirabilis, Salmonella species, Staphylococci and Streptococci in cattle, sheep, dogs and cats.

### INDICATIONS:

Selmox LA injection is indicated for the treatment of respiratory, gastrointestinal, urogenital tract infections, skin and soft tissues infections and in prevention of post-operative infections. It is a drug of choice for the treatment of pneumonia, ear and throat infections, enteritis, E.coli, wound infections, mastitis, and metritis. In poultry, for coryza, enteritis, streptococcosis and other respiratory disease.

### DOSAGE AND ADMINISTRATION:

Administer the following dose by intramuscular or subcutaneous injection:

**Cattle / Buffalo / Sheep / Goat:** 1ml per 13.2kg body weight. Repeat the dose after 48 hours if required.

**Poultry:** 0.2ml per kg body weight.

### CONTRA INDICATIONS:

This product is not suitable for administration via intravenous or intrathecal routes.

Milk for human consumption must not be taken from a cow during treatment. Milk for human consumption may only be taken after 24 hours from the last treatment.

Not for use in sheep producing milk for human consumption.

Animals must not be slaughtered for human consumption during treatment. Cattle may be slaughtered for human consumption only after 23 days from the last treatment. Sheep may be slaughtered for human consumption only after 16 days from the last treatment.

Occasional local tissue reaction may result from use of this product.

As with other penicillins Amoxicillin should not be used in rabbits, hamsters.

### OPERATOR WARNINGS:

Penicillin and cephalosporins may cause hypersensitivity (allergy) following injection, inhalation, ingestion or skin contact. Hypersensitivity to penicillin may lead to cross reactions to cephalosporins and vice versa. Allergic reactions to these substances may occasionally be serious. Do not handle this product if you have known sensitivity to penicillins or if you have been advised not to work with such preparations.

Handle this product with great care to avoid exposure taking all recommended precautions.

If you develop symptoms following exposure, such as a skin rash, you should seek medical advice and show the doctor this warning. Swelling of the face, lips or eyes or difficulty in breathing are more serious symptoms and require urgent medical attention.

### PRECAUTIONS:

**Shake well before use.**

Do not use in animals sensitive to penicillin.

Consult the veterinarian before use.

Keep out of the reach of children.

Store between 15-25°C in a cool and dry place.

### WITHDRAWAL TIME:

**Meat:** 28 days.

**Milk:** 03 days.

BP Vet. Specs.



Manufactured by:

**Selmore**

Pharmaceuticals (Pvt) Ltd.

36-Km Multan Road, Lahore Pakistan.



ISO CERTIFIED COMPANY