

TYGENT

Injection

For Veterinary Use Only

STATEMENT OF ACTIVE SUBSTANCE (S) AND OTHER INGREDIENTS:

Each ml contains

Tylosin Tartrate.....100mg
Gentamicin (as Sulphate).....50mg

INDICATIONS:

Tygent Injection is indicated in respiratory tract infections, gastrointestinal tract infections and genital tract infections like CRD, pneumonia, bronchitis, pleuritis, pasteurellosis, enteritis, gastroenteritis, salmonellosis, metritis and mastitis etc.

CONTRAINDICATIONS:

Do not use in animals with known hypersensitivity to the active ingredient. Do not use in animals with impaired renal function. Do not use in pregnant animals. Do not administer to horses.

ADVERSE REACTIONS:

Localised, painless, oedematous swelling may occasionally be seen at the injection site.

TARGET SPECIES:

Cattle, sheep, goat

DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION:

For intramuscular injection. Large Animals: 1ml per 25-30kg body weight.

Sheep / Goat: 1ml per 20kg body weight.

ADVICE ON CORRECT ADMINISTRATION:

None.

WITHDRAWAL PERIOD (S):

Meat and offal: 30 days Milk: 7 days

PRECAUTIONS:

Keep out of the sight and reach of children. Store between 15-25°C in a cool and dry place.

Do not use this veterinary medicinal product after the expiry date which is stated on the carton.

SPECIAL WARNINGS INCLUDING:

Special warnings for each target species:

None.

Special precautions for use in animals:

Tygent must be administered by intramuscular route. Do not exceed the stated dose.

Special precautions to be taken by the person administering the veterinary medicinal product to animals: Wash hands before and after treatment.

Pregnancy and lactation:

The use during pregnancy is contra-indicated due to the possible

nephro- and ototoxicity in the foetus. Interaction with other medicinal products and other forms of interaction: Local swelling at the injection site can occur.

Quickly acting diuretics administered concurrently with gentamicin increase the likelihood of nephrotoxicity and the ototoxicity.

Gentamicin potentiates the action of the general anaesthetics and muscle relaxants resulting in a higher risk of neuromuscular block. Halothanes raise the cardiovascular depressing effect of gentamicin. Overdose (symptoms, emergency procedures, antidotes): The therapeutic index of tylosin is very large. The risk of overdose is therefore very small.

Relative overdosing resulting in neuromuscular block is possible when gentamicin is administered concurrently with general anaesthetics and/or muscle relaxants.

Incompatibilities:

In the absence of compatibility studies, the veterinary medicinal product must not be mixed with other veterinary medicinal products.


SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS IF ANY:

Medicines should not be disposed of via wastewater or household waste. Ask your veterinarian how to dispose of medicines no longer required. These measures should help to protect the environment.

DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED:

21-12-2020.

Selmore's Specs



Manufactured by:
Selmore Pharmaceuticals (Pvt) Ltd.
36 Km Multan Road, Lahore, Pakistan.

