# ZEPTOCOL

For Veterinary Use Only

### Composition:

Each kg powder contains:

Procaine Penicillin	.12gm
Streptomycin Sulphate	
Colistin Sulphate	
Zinc Bacitracin	52gm

#### Description:

Zinc bacitracin was effective as a growth promoter in the broilers reared during 1 to 21 days. In the study period, the use of zinc bacitracin resulted in lower feed intake and greater weight gain, when compared to the treatment without antibiotic Colistin a Cyclic Polypeptide antibiotic has a bactericidal selected action against gram-negative bacteria, the primary bacterial effect of colistin is a blockade of the cell division of bacteria followed by a secondary progressive lyses.

## Indications:

Zeptocol Powder is indicated for the control and prevention of respiratory infections, CRD, colibacillosis, salmonellosis and early chick mortality, increase in egg production, hatchability and immune system for the increase in production as a growth promoter.

#### Dosage and Administration:

Increase weight gain and improve FCR: 500g-1kg per ton of feed continuously. Increase of egg production and hatchability: 1kg-2kg per ton of feed continuously.

Prevention and treatment of disease during stress period: 2.5kg to 5kg per ton of feed continuously.

#### Precautions:

Do not use in animals sensitive to penicillin. aminoglycosides & colistin. Following withdrawal of the first dose, use the product within 28 days. Consult the veterinarian before use. Protect from light & moisture. Keep out of the reach of children.

Store between 15-25°C in a cool and dry

Innovator's Specs.

place.





